

Care guide

Wuelfing's Stick Insect, *Acrophylla wuelfingi*



Wuelfing's Stick Insects are extremely large straight bodied insects, and are among Australia's longest species. They have an array of small knobs or spines on their thorax, and have large serrations on the upper legs. Like most phasmids, the females are much larger and more robust than the males, and can grow to a length of 210mm. Mature females have large fan-like wings which are used to startle predators, but cannot fly due to their body weight. Mature males are light-weight and can fly quite well.

They have a very interesting reproductive cycle, beginning with the eggs being tossed individually by females from the trees down to the forest floor. A single female may lay hundreds of eggs in a lifetime. The eggs are matte black and look very much like plant seeds. They are often collected by ants and stored below ground in their nests which protects the eggs from predators.

After hatching, the baby stick insects (nymphs) must make their way to the surface and into a tree. The nymphs are green and slender at this stage, and use their appearance to blend into the small growth at the tips of branches.

Once in a tree they begin to feed on leaves and grow by shedding their outer skeleton (exoskeleton). This process is called ecdysis or moulting. To moult successfully the insect needs to hang uninterrupted beneath a leaf or branch. This can take 10mins to half an hour. Both sexes change from bright green to olive brown as they approach maturity, although individuals may go through various multi-coloured combinations along the way.

An amazing fact about many phasmids is that they are parthenogenetic. This means that females don't need to be mated to reproduce. So if you only have a single female, she can still produce young - replicas of herself.

Care guide

Wuelfing's Stick Insect, *Acrophylla wuelfingi*

Food: Various species of Eucalyptus (gum), Acacia (wattle), Cadagi and Guava. Ensure fresh, healthy leaves are always available to your insect. Don't let the leaves dry out before you change them. The leaves can be put into a jar of water to keep them fresh for as long as possible. Make sure the jar has a lid or covering with holes in it to stop the young stick insects from falling in and drowning. *Offering two or three species of food plant when you first get your stick insects is a good way finding one that they like.

Water: Mist-spray the leaves around your insect once a day – it will drink the droplets.

Enclosure: The enclosure needs to be large enough to allow your stick insect to shed its exoskeleton properly. A number of individuals can be housed together as long as each insect has sufficient space to hang and feed without disturbing others. An enclosure should be higher than it is wide, as stick insects like to climb upwards. Place the enclosure in a spot where it gets a bit of daylight each day, but be careful it doesn't over heat in direct sunlight. Make sure there is plenty of ventilation in your enclosure.

Temperature: This species will do best at 22 – 28° C, but will tolerate 10 – 20° C and have a slower growth rate.

Life span: 6 months – 1 year

Handling: They may be picked up gently, by coaxing them onto your hand. They always prefer to climb up, so use this to your advantage when picking them up and putting them back onto the leaves. They do have small hooks on their feet that they use to hang on with – these may tickle your hand a little.

Reproduction: Adult males will mate with adult females all year round. Females may also lay eggs even when she hasn't been mated. Once the female starts to lay eggs, they can be collected from the bottom of the enclosure and stored in a small container. From 3 -9 months, the eggs may start hatching. Incubation times will be shorter at warmer temperatures, and are best incubated at 23- 26 degrees. The hatchlings need to be fed on soft, young leaves, and can be housed in the same enclosure as the adults.

Common issues: Sometimes stick insects will not shed their exoskeleton properly. This may be because there isn't enough room in the enclosure for them to do so, or that their environment is too dry and their new exoskeleton hardens before they can free themselves from the old one. They can also lose limbs through the moulting process if they aren't hanging correctly whilst moulting. They can regrow lost limbs at their next moult, but need at least two moults to regrow a limb to a usable size.

NOTE: These animals are captive bred, and should not be released into the wild.

For more information contact Minibeast Wildlife at info@minibeastwildlife.com.au